



ISSN Print: 2394-7489
ISSN Online: 2394-7497
IJADS 2020; 6(2): 628-630
© 2020 IJADS
www.oraljournal.com
Received: 01-02-2020
Accepted: 03-03-2020

Zuhair Al-Nerabiah
Pediatric Dentistry Department,
Dental Collage, Damascus
University, Syria

Muaaz Alkhouli
Pediatric dentistry Department,
Dental Collage, Damascus
University, Syria

Mohannad Laflouf
Professor, Pediatric Dentistry
Department, Dental Collage,
Damascus University, Syria

Mahmoud Abdul-Hak
Professor, Oral Medicine
Department, Dental Collage,
Damascus University, Syria

Corresponding Author:
Zuhair Al-Nerabiah
Pediatric Dentistry Department,
Dental Collage, Damascus
University, Syria

Pediatric dentists consideration for Covid-19 in children: Review article

Zuhair Al-Nerabiah, Muaaz Alkhouli, Mohannad Laflouf and Mahmoud Abdul-Hak

Abstract

As of June 2020, over 6 million confirmed cases of Covid-19 (Corona virus) infection and nearly 400 thousand Covid-19-related deaths have been reported by the World Health Organization globally. Covid-19 may occur sporadically in communities or may be transmitted within families or hospitals. The number of confirmed Covid-19 cases among healthcare workers has been increasing. Covid-19 may also spread through aerosols generated during various dental treatments, resulting in transmission between patients and dentists. As Covid-19 cases have also been reported among children, pediatric dentists are at risk of Covid-19 infection. This review discusses Covid-19 infection in children and healthcare workers, especially pediatric dentists, and considerations pertaining to pediatric dentistry since the risk of Covid-19 transmission from an infected patient may be high due to the unique work environment of dentists (aerosol generation).

Keywords: Covid-19, children, pediatric dentist

Introduction

The coronavirus disease (COVID-19) is a novel viral infection that started in Wuhan, China and caused the outbreak of pneumonia in the rest of the world. It seems that the rapidly spreading virus is more contagious than severe acute respiratory syndrome coronavirus and Middle East respiratory syndrome coronavirus [1].

Covid-19 is a novel beta coronavirus of the Coronaviridae family that causes a severe respiratory disease with a high fatality rate [2].

As of June 2020, over 6 million confirmed cases of Covid-19 (Corona virus) infection and nearly 400 thousand Covid-19-related deaths have been reported by the World Health Organization globally. The incubation period for human to human transmission ranges from 2-15 days [2].

The symptoms of Covid-19 infection range from being asymptomatic to severe pneumonia, acute respiratory distress syndrome, septic shock, and multi-organ failure, leading to death. The symptomatic patients may have fever, cough, chills, throat soreness, myalgia, arthralgia, vomiting, or diarrhea. Furthermore, men over 45 years of age, patients with comorbidities, and healthcare workers have been reported to be high-risk groups for Covid-19 infection [3].

The virus has been detected in respiratory, gastrointestinal and other bodily secretions, as well as in air samples, which indicates the possibility for airborne transmission [4].

Several studies have reported Covid-19 infections among healthcare workers and children. Viral infections, such as severe acute respiratory syndrome, may be transmitted to healthcare workers from infected patients through aerosols. Considering that several types of dental equipment, such as handpieces, air-water syringes and ultrasonic scalers, produce considerable amounts of aerosols, the potential for the spread of infections from patients to dentists or dental assistants is high [5].

A suggested route of human-to-human transmission is through airborne droplets, touching or coming into contact with an infected person or a contaminated surface. Moreover, other routes such as blood or saliva have not been explored but are possible because of the documented transmission of blood-borne infectious diseases such as HIV/AIDS, hepatitis C virus, and hepatitis B virus through blood or saliva. These routes of transmission increase the concern about a similar route of transmission for COVID-19 in the dental setting [6].

This review is an attempt to discuss Covid-19 infection among children and those providing dental treatment to them, including precautions and considerations pertaining to the practice of pediatric dentistry.

Covid-19 infection in healthcare workers

The 3 patterns of Covid-19 transmission suggested in the literature include the following: a) sporadic cases occurring in communities; b) transmission within families; and c) nosocomial transmission. Health care related Covid-19 transmission is associated with high morbidity, extended use of mechanical ventilation and fatality rates of up to 5%; this type of transmission can be attributed to shortcomings in observing stringent infection control protocols [7].

Furthermore, Covid-19 has been reported to be viable in hospital like environments for up to 48 hours with a stability that is unaltered during aerosolization [8]. Healthcare workers could be infected with Covid-19 through exposure in the community or at their workplace, they could be diagnosed late, and they could remain asymptomatic or mildly symptomatic [9].

Furthermore, unsuspected cases entering healthcare facilities have been considered the main source of MERS-CoV.4 Considering these aspects and that healthcare workers may continue to work regardless of being symptomatic, the possibility of transmitting the infection to their patients is also high [10].

Covid-19 infection in children

Reports have concluded that although Covid-19 infection presents with a wide range of clinical manifestations, the mortality rate in children is lower than that in adults. A review on the effects of Coronavirus infection in children concluded that Human Coronavirus infections may be associated with respiratory symptoms and may also involve central nervous system. The authors suggested that the clinical and genetic traits of Human Coronavirus infection among children should be monitored closely for early prevention [11].

Dental considerations

Bioaerosols in dental practice. Bioaerosols are defined as a suspension of biological particles in gaseous media. Apart from radiation exposures, dermatitis, eye injuries, and musculoskeletal and respiratory disorders, other occupational health risks exist in dentistry. These risks include percutaneous exposure incidents and exposure to infectious diseases, such as those that may be present in bioaerosols. Subgingival scaling of periodontally involved teeth with ultrasonic scalers may produce aerosols containing blood [12].

One study reported that ultrasonic scalers and tips produced significantly more aerosol and splatter than a handheld curette, regardless of the scaler type used [13]. Recent report studied the characteristics of blood-containing aerosols generated by common powered dental instruments. The author concluded that all the recovered particles could contain the hepatitis B virus and be inhaled and retained (20-100%) in the human respiratory system. The repeated and chronic exposure to bioaerosols generated during certain dental procedures and the relatively small particle size contribute to an increased risk of infection among dental professionals [13].

Furthermore, the protection provided by surgical masks worn by dental professionals may be low for small particles; in addition, these masks may not fit perfectly in clinical practice [14].

A study reported that 15-83% of plasma aerosol particles ranging from 0.06-2.5 microns in size passed through the

filters of 9 makes of surgical masks used by dentists. These factors may play a role in the airborne route of viral infections among dentists and clinical dental auxiliaries. The likelihood of detecting, reporting, documenting and publishing dentistry-associated infections is relatively less [15].

Although no definitive evidence of an extensive public health hazard from exposure to dental unit waterlines has been reported. Few case reports have suggested a definitive link between exposure to contaminated dental unit waterlines and Legionella infection [16].

Infection control in dental practice

A recent review on the emerging and fluctuating viral diseases in the new millennium concluded that infection control plays an equally important role in the practice of dentistry as do the understanding of oral manifestations and the diagnosis and management of viral infections [17].

While differences exist in the virology, epidemiology, and clinical outcomes of Covid-19 and SARS-CoV infections, the clinical symptoms of Covid-19 resemble those of SARS-CoV, apart from acute renal failure [2].

Although these results cannot be directly interpreted for a definitive conclusion, the management protocols for different Covid-19 infection-associated with clinical scenarios for dental professionals may be similar to those of SARS-CoV.

The implications of SARS-CoV for general dental practitioners, the significance of droplets and aerosols in disease transmission, the recommended management protocols for SARS-CoV infection and specific infection control measures have been well described by Li *et al.*

A comprehensive review on universal and standard precautions has been published elsewhere and is beyond the scope of this article, infection control measures pertaining only to pediatric dental practice in the context of Covid-19 infection are discussed here [14].

In pediatric dental practice, effective infection control measures for the prevention or minimization of viral infection transmission can be implemented by a) controlling the gag or cough reflex; b) reducing aerosol/ splatter generation; c) managing contaminated air and; d) improving personal protection.

The gag or cough reflex may be stimulated by certain procedures, such as posterior intraoral and bite-wing radiographs and taking impressions.

Orthopantomographs or oblique lateral views may be considered instead of intraoral radiographs for screening, whereas oral mucosa in very sensitive patients may be anesthetized before taking impressions. Sedation may also be considered to control gag reflex. Varying levels of physical, intellectual, emotional and social development of children and adolescents are major challenges for dentists especially attending to their psychological needs within adequate infection control regimen.

Toys provided for pediatric patients may be a potential source of cross-infection. Soft toys are more likely to be contaminated, difficult to disinfect and may re-contaminate quickly compared to hard surfaced toys. Furthermore, restraining devices used to control movements of pediatric patients such as Velcro fasteners may also be contaminated and should be disinfected accordingly [18].

Extra-oral evacuation devices and special aerosol reduction devices may be used in combination with ultrasonic scalers to reduce the amount of aerosol produced.

In high-risk cases, chemo-mechanical caries removal or the atraumatic restorative technique may be utilized to prevent

or reduce aerosol and splatter generation. In addition, high-volume evacuation removes infectious droplets at the source as they are emitted; thereby, minimizing, or preventing their dispersion in the air^[19].

To maintain their efficacy, the filters in the suction apparatus should be cleaned every day, and the exhaust air should be vented outside to prevent the recirculation of contaminated air. Contaminated air can be managed by improving dental clinic ventilation and/or by disinfecting the air. An ideal airflow pattern combined with a minimum of 3 air changes per hour has been recommended for dental clinics^[20].

Moreover, although its use in dental clinics is unconfirmed, ultraviolet germicidal irradiation may be installed and is effective against fungi, viruses, and bacteria, namely, tubercle bacilli and anthrax. On the other hand, measures of improving personal protection include washing hands frequently before and after treatment, using disposable barriers, dispensing instruments and materials just before treatment (thereby preventing particles from settling on the surfaces), and sterilizing soiled instruments^[20].

After each patient visit, surfaces may be disinfected using hospital-grade disinfectants, which are effective against coronavirus. Personal protective equipment, such as gowns, hair covers, masks, gloves, shielded face masks and shoe covers, should be used as appropriate.

A higher level of respiratory protection should be considered, especially with aerosol-generating procedures. The use of N95 respirators offers a certain level of protection against the airborne transmission of SARS-CoV-2, although the exact level of protection offered by these respirators for individuals may vary.

These respirators may also provide some level of protection against Covid-19 and may be utilized instead of surgical masks in dental clinics. Apart from these measures, vaccination of pediatric dentists and staffs working in pediatric dental clinics against measles, mumps, varicella and rubella may be necessary for their protection from these viral infections^[21].

Conclusion

In conclusions, considering the unique work environment of dentists, which involves close patient contact and aerosol production, the risk of Covid-19 transmission from an infected patient is high. As the number of Covid-19 cases may increase in future, pediatric dentists should be well informed and educated about not only the signs and symptoms of the condition but also how to follow stringent infection control measures in these cases.

Conflict of interest

Authors declare they have no conflict of interest

Acknowledgment

This study was funded by the University of Damascus.

References

1. COVID CDC, Team R. Severe outcomes among patients with coronavirus disease 2019 (COVID-19)-United States, February 12–March 16, 2020. *MMWR Morb Mortal Wkly Rep.* 2020; 69(12):343-6.
2. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun.* 2020; 102433.
3. Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, evaluation and treatment coronavirus

(COVID-19). In: Statpearls [internet]. StatPearls Publishing, 2020.

4. Novel CPERE. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi.* 2020; 41(2):145.
5. Meng L, Hua F, Bian Z. Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res.* 2020; 99(5):481-7.
6. Spagnuolo G, De Vito D, Rengo S, Tatullo M. COVID-19 outbreak: an overview on dentistry. *Multidisciplinary Digital Publishing Institute,* 2020.
7. Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S *et al.* Early dynamics of transmission and control of COVID-19: A mathematical modelling study. *lancet Infect Dis,* 2020.
8. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W *et al.* Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet.* 2020; 395(10226):809-15.
9. Guo YR, Cao QD, Hong ZS, Tan YY, Chen SD, Jin HJ, *et al.* The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Mil Med Res.* 2020; 7(1):1-10.
10. Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, Wang M. Presumed asymptomatic carrier transmission of COVID-19. *Jama.* 2020; 323(14):1406-7.
11. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z *et al.* Epidemiology of COVID-19 among children in China. *Pediatrics,* 2020, 145(6).
12. Sabino-Silva R, Jardim ACG, Siqueira WL. Coronavirus COVID-19 impacts to dentistry and potential salivary diagnosis. *Clin Oral Investig.* 2020, 1-3.
13. Ge Z, Yang L, Xia J, Fu X, Zhang Y. Possible aerosol transmission of COVID-19 and special precautions in dentistry. *J Zhejiang Univ B.* 2020, 1-8.
14. Coulthard P. Dentistry and coronavirus (COVID-19)-moral decision-making. *Br Dent J.* 2020; 228(7):503-5.
15. Bartoszko JJ, Farooqi MAM, Alhazzani W, Loeb M. Medical masks vs N95 respirators for preventing COVID-19 in healthcare workers: A systematic review and meta-analysis of randomized trials. *Influenza Other Respi Viruses,* 2020.
16. Barbeau J, Gauthier C, Payment P. Biofilms, infectious agents, and dental unit waterlines: a review. *Can J Microbiol.* 1998; 44(11):1019-28.
17. Cheng VCC, Wong SC, Chen JHK, Yip CCY, Chuang VWM, Tsang OTY *et al.* Escalating infection control response to the rapidly evolving epidemiology of the Coronavirus disease 2019 (COVID-19) due to SARS-CoV-2 in Hong Kong. *Infect Control Hosp Epidemiol.* 2020; 41(5):493-8.
18. Izzetti R, Nisi M, Gabriele M, Graziani F. COVID-19 transmission in dental practice: brief review of preventive measures in Italy. *J Dent Res.* 2020, 0022034520920580.
19. Alharbi A, Alharbi S, Alqaidi S. Guidelines for dental care provision during the COVID-19 pandemic. *Saudi Dent J,* 2020.
20. Carmona JR. COVID-19 Outbreak and Dentistry. *Odovtos-International J Dent Sci.* 2020; 22(2):9.
21. Ren YF, Rasubala L, Malmstrom H, Eliav E. Dental care and oral health under the clouds of COVID-19. *JDR Clin Transl Res.* 2020, 2380084420924385.