



## International Journal of Applied Dental Sciences

ISSN Print: 2394-7489  
ISSN Online: 2394-7497  
IJADS 2017; 3(2): 193-196  
© 2017 IJADS  
www.oraljournal.com  
Received: 02-02-2017  
Accepted: 03-03-2017

**Subija K Narayanan Kutty**  
MDS, Consultant Endodontist,  
Thrissur, Kerala, India

**MS Lekshmi**  
BDS, Junior Resident,  
Department of Conservative  
Dentistry and Endodontics,  
Government Dental College,  
Thiruvananthapuram, Kerala,  
India

**Aparna Mohan E**  
BDS, Junior Resident,  
Department of Conservative  
Dentistry and Endodontics,  
Government Dental College,  
Thiruvananthapuram, Kerala,  
India

**Lija Isaac**  
BDS, Junior Resident,  
Department of Conservative  
Dentistry and Endodontics,  
Government Dental College,  
Thiruvananthapuram, Kerala,  
India

**Correspondence**  
**Subija K Narayanan Kutty**  
MDS, Consultant Endodontist,  
Thrissur, Kerala, India

### Pulp tissue dissolution in endodontics- A review

**Subija K Narayanan Kutty, MS Lekshmi, Aparna Mohan E and Lija Isaac**

#### Abstract

Meticulous shaping and cleaning of the root canals is of utmost importance to the success of endodontic therapy. A combination of mechanical instrumentation with appropriate chemicals is conventionally used for thorough cleaning and disinfection of the complex root canal system. Ideally, an irrigant should be antimicrobial, biocompatible, dissolve pulp tissue, mechanically flush debris and help to remove smear layer. The capacity to dissolve pulp tissue, both vital and necrotic, is of paramount importance, as the solution can help clean areas of the root canal that are inaccessible to mechanical instrumentation. This paper reviews various irrigants with respect to their pulp dissolution capacity.

**Keywords:** Pulp tissue, dissolution, irrigant

#### Introduction

The main objectives of endodontic treatment are thorough shaping and cleaning of the pulp spaces followed by complete obturation with an inert material. The root canal system is complex, and canals may branch and divide. Variations in anatomy such as fins, isthmi and C-shaped canals may be present [1]. Studies have shown that approximately 35-40% of the canal walls remain uninstrumented because of these complexities [2]. Geometrically symmetrical instruments used for shaping root canals will not be able to reach these anatomically complex, inaccessible areas [3]. Therefore, there is a need to supplement mechanical preparation with chemicals which can dissolve the tissue remnants and disinfect the canal system. The mechanical action of instruments and chemical effect of irrigants occur simultaneously, referred to as chemomechanical preparation. Ideally, an endodontic irrigant must be antimicrobial, dissolve pulp tissue remnants and remove the smear layer. It should be non-irritating to the periapical tissues, non-antigenic, non-toxic, and non-carcinogenic [4]. Pulp tissue remnants may provide a source of nutrition for surviving bacteria, may cause severe pain, and may interfere with proper obturation, thereby leading to endodontic treatment failure [5]. Therefore, tissue dissolution is one of the most important properties that are required of an endodontic irrigant.

#### Methods used to assess tissue dissolution *in vitro*

Different methods have been used to measure the dissolution process *in vitro*- analysis of the hydroxyproline content in the residual tissue (a measure of dissolution of collagen in the tissue), analysis of hydroxyproline in the irrigant, measurement of available chlorine in the solution after dissolution process, assay of radioactive iodine in the irrigant, scanning electron microscopy, measurement of weight loss of tissue over a specified time, and assessment of the time needed for complete dissolution of the specimen [6-8]. Of these, measurement of weight change and time to complete dissolution are the most widely accepted and used methods. However, it is difficult to ascertain the end-point of complete dissolution as a large number of bubbles are formed due to saponification reaction as in the case of irrigant like sodium hypochlorite [6, 7]. Measuring the weight change in a particular time interval is more relevant to the clinical scenario as compared to evaluating the time taken for complete dissolution.

#### Commonly used and potential irrigants investigated for organic tissue dissolution capacity

##### Sodium hypochlorite

Sodium hypochlorite (NaOCl) is an excellent antimicrobial and tissue dissolving agent. It is the most widely used endodontic irrigant, at concentrations ranging from 0.5-6%.

This chemical has been extensively investigated with respect to its mechanism of action and factors which affect its properties [1]. However it is extremely toxic. Contact of NaOCl with organic matter like dentin and pulp results in the formation of organochlorine compounds which may be carcinogenic [9].

### Mechanism of action of NaOCl

Estrela *et al.* [10] described the mechanism of action of NaOCl with respect to its tissue dissolving and antimicrobial properties. NaOCl in aqueous medium exhibits a dynamic balance as shown by the reaction:



Three types of reactions take place due to the interaction of NaOCl with organic matter. These reactions take place mostly at the surface, and occur simultaneously and synergistically.

**Saponification reaction-** NaOCl functions as an organic and fat solvent, degrading fatty acids and converting them into fatty acid salts (soap) and glycerol (alcohol), and reduces surface tension of the remaining solution [10].

**Amino acid neutralization reaction-** NaOCl reacts with amino acids to form salt and water, with a reduction of pH and exit of hydroxyl ions [10].

**Chloramination reaction-** Hypochlorous acid (HOCl) present in NaOCl solution is an organic tissue solvent. It releases chlorine, which form chloramines on reaction with amino group of proteins. Hypochlorous acid and hypochlorite ions (OCl<sup>-</sup>) cause amino acid degradation and hydrolysis [10].

### Factors affecting tissue dissolution

Organic tissue dissolution is significantly influenced by tissue type, concentration and pH of the solution, time of exposure and rate of replenishment, temperature, amount of organic matter relative to the volume of solution, frequency and intensity of agitation, available surface area of tissue, canal preparation size, and mechanical action [11, 12].

#### i. Type of tissue

A vast variety of tissues have been used in studies assessing tissue dissolution - porcine muscle, porcine palatal mucosa, porcine pulp, bovine muscle, bovine pulp, human pulp, rabbit liver, and rat connective tissue [7]. Human pulp tissue has been used only in a few studies. Tissues other than pulp have been used due to their easy availability and the relative ease to standardize the various parameters that affect dissolution such as surface area, volume and weight of tissue [13]. However, a study revealed that NaOCl solution took approximately three times longer to dissolve porcine palatal mucosa completely when compared to human pulp [14].

#### ii. Concentration and pH of the solution

Higher concentrations of NaOCl produce greater dissolution of organic tissue [11]. Higher bactericidal activity is demonstrated by solutions with lower pH. However tissue dissolving activity is enhanced with increase in pH and concentration [12]. This can be explained by the fact that higher the concentration and pH, greater is the number of NaOH molecules, which play a considerable role in tissue dissolution. After dissolution, pH decreases, and the dynamic equilibrium is shifted towards increased formation of HOCl and decreased dissolution velocity [15].

#### iii. Time of exposure and rate of replenishment of solution

Initial reaction between NaOCl and organic matter occurs relatively fast, followed by a slower phase of dissolution. The reaction occurs mainly on surface of the tissue, and molecules of NaOCl are consumed in the reaction, leading to their local depletion. Most of the activity of NaOCl is lost after two minutes of contact with organic tissue. Frequent replenishment of the solution is therefore essential to ensure availability of active molecules for further reaction and to remove remnants of dissolved tissue [7, 11].

#### iv. Temperature

Dissolution capacity of NaOCl increases with increase in temperature [16, 17].

#### v. Amount of organic matter in relation to volume of solution

For optimum activity of an irrigant, volume of the solution should be in excess

compared to the amount of tissue to be dissolved. Excess organic matter depletes available chlorine in NaOCl and causes pH to drop, thereby decreasing its activity [7, 11]. Therefore, continual replenishment with fresh irrigant is essential during instrumentation.

#### vi. Frequency and intensity of agitation

Mechanical agitation is an important factor in promoting tissue dissolution, as it influences the fluid flow [1].

#### vii. Surface area in contact with the solution

Dissolution mainly occurs on the surface of tissue in contact with the solution. In

the clinical situation, the irrigant comes in contact with only a small portion of the pulp tissue, unlike in studies conducted *in vitro*, where the tissue is usually immersed in the solution. Several studies have used bovine and porcine muscle, or porcine palatal mucosa, in order to obtain samples of uniform surface areas. Human pulp tissues present a major challenge to obtaining uniform samples.

#### viii. Surface tension

In less concentrated NaOCl solutions, there is greater interaction of HOCl with organic matter. In more concentrated solutions, there is greater interaction of NaOH, leading to greater reduction of surface tension. Higher the initial concentration of NaOCl, greater is the surface tension reduction. This is due to the fact that solutions with higher active chlorine concentration, and therefore higher NaOH concentration, provide greater formation of soaps, which reduce the surface tension. Greater dissolution capacity of NaOCl has been noted in the presence of surfactants. This may be due to the better contact of the solution with the tissue [18, 19].

#### ix. Stability of NaOCl

Solutions of NaOCl are unstable. The amount of available chlorine deteriorates with time, exposure to heat and light, and on contact with air, metals, metallic ions and organic matter. To ensure good shelf life, the solution should be stored in light-proof opaque glass or polythene airtight containers, in a cool place [4].

#### x. Effect of dentin

Dentin is found to have a detrimental effect on tissue dissolution potential of NaOCl [20, 21].

#### xi. Effect of other irrigants

Presence of EDTA and hydrogen peroxide adversely affected dissolution by NaOCl [21, 22].

#### xii. Ultrasonic activation

Passive ultrasonic irrigation promotes tissue dissolving effects of NaOCl [23].

#### xiii. Effect of lasers

The effect of Er:YAG laser irradiation has been investigated and it was concluded that laser activation of NaOCl at 200 mW output led to effective dissolution [24].

#### xiv. PIPS

Photon-initiated photoacoustic streaming technique of irrigant activation was found to increase the dissolution potential of NaOCl [25].

#### xv. Ozone

HealOzone was found to increase speed of dissolution by NaOCl [26].

#### Calcium hydroxide

Calcium hydroxide has minimal pulp dissolution potential. Calcium hydroxide paste causes better dissolution of necrotic remnants, but not the solution [13, 27].

#### Hydrogen peroxide

Combination of hydrogen peroxide with NaOCl was found to delay dissolution by NaOCl [22].

#### Chlorhexidine

Chlorhexidine, though an excellent antibacterial, is ineffective as a solvent of pulp tissue [14, 28, 29]. However 2% chlorhexidine gel associated with 8% papain gel was found to dissolve pulp tissue [30].

#### Ethylene diamine tetraacetic acid (EDTA)

EDTA exhibits minimal dissolution of organic tissue [31, 32]. Furthermore it adversely affects the dissolution potential of NaOCl [33].

#### MTAD

MTAD is not an effective solvent of organic tissue [29].

#### Chlorine dioxide

Different concentrations of chlorine dioxide have shown ability to dissolve pulp tissue, but to a lesser extent than NaOCl [34-36].

#### Peracetic acid

Pulp dissolution has been reported with 5% peracetic acid, though significantly less than NaOCl [37].

#### Papain gel

8% papain gel was found to dissolve pulp tissue, both alone and in combination with chlorhexidine gel [30].

#### Calcium hypochlorite

5% and 10% solutions of calcium hypochlorite were assessed and found to be good solvents of pulp tissue [36, 38].

#### Maleic acid

7% maleic acid was found to have minimal dissolution potential when compared to NaOCl [32].

Apart from these, several other chemicals have been tested for

their tissue dissolving abilities. Citric acid [14], dichloroisocyanurate [14], super-oxidized water [26], octenidine [39], garlic extract [40] etc. have been investigated but found to be ineffective.

#### Conclusion

Over the years, several chemicals and combinations have been investigated for potential use as endodontic irrigant. Sodium hypochlorite, though toxic, is the most commonly used irrigant as it is an excellent antimicrobial agent and solvent of organic tissue. Despite extensive research, an ideal irrigant with all the required properties is yet to be identified. Some chemicals have shown optimistic results as far as pulp dissolution is concerned. Further research should be directed towards understanding their mechanism of action on pulp tissue and thereby the various methods which would enhance pulp dissolution.

#### References

- Hargreaves KM, Cohen S, Berman LH. Cohen's Pathways of the Pulp. Mosby Elsevier, 2011.
- Peters OA, Schönenberger K, Laib A. Effects of four Ni-Ti preparation techniques on root canal geometry assessed by micro computed tomography. *Int Endod J*. 2001; 34(3):221-30.
- Sum CP, Neo J, Kishen A. What we leave behind in root canals after endodontic treatment: some issues and concerns. *Aust Endod J Aust Soc Endodontology Inc*. 2005; 31(3):94-100.
- Basrani B, Haapasalo M. Update on endodontic irrigating solutions. *Endod Top*, 2012; 27(1):74-102.
- Ingle JI, Bakland LK, Baumgartner JC. Ingle's Endodontics. PMPH-USA, 2008.
- Clarkson R, Moule A, Podlich H, Kellaway R, Macfarlane R, Lewis D *et al*. Dissolution of porcine incisor pulps in sodium hypochlorite solutions of varying compositions and concentrations. *Aust Dent J*. 2006; 51(3):245-51.
- Stojicic S, Zivkovic S, Qian W, Zhang H, Haapasalo M. Tissue dissolution by sodium hypochlorite: effect of concentration, temperature, agitation, and surfactant. *J Endod*. 2010; 36(9):1558-62.
- Wadachi R, Araki K, Suda H. Effect of calcium hydroxide on the dissolution of soft tissue on the root canal wall. *J Endod*. 1998; 24(5):326-30.
- Varise TG, Estrela C, Guedes DFC, Sousa-Neto MD, Pécora JD. Detection of Organochlorine Compounds Formed During the Contact of Sodium Hypochlorite with Dentin and Dental Pulp. *Braz Dent J*. 2014; 25(2):109-16.
- Estrela C, Estrela CRA, Barbin EL, Spanó JCE, Marchesan MA, Pécora JD. Mechanism of action of sodium hypochlorite. *Braz Dent J*. 2002; 13(2):113-7.
- Moorer WR, Wesselink PR. Factors promoting the tissue dissolving capability of sodium hypochlorite. *Int Endod J*. 1982; 15(4):187-96.
- Christensen CE, McNeal SF, Eleazer P. Effect of lowering the pH of sodium hypochlorite on dissolving tissue *in vitro*. *J Endod*. 2008; 34(4):449-52.
- Türkün M, Cengiz T. The effects of sodium hypochlorite and calcium hydroxide on tissue dissolution and root canal cleanliness. *Int Endod J*. 1997; 30(5):335-42.
- Naenni N, Thoma K, Zehnder M. Soft Tissue Dissolution Capacity of Currently Used and Potential Endodontic Irrigants. *J Endod*. 2004; 30(11):785-7.
- Palazzi F, Morra M, Mohammadi Z, Grandini S,

- Giardino L. Comparison of the surface tension of 5.25% sodium hypochlorite solution with three new sodium hypochlorite-based endodontic irrigants: Surface tension of endodontic irrigants. *Int Endod J.* 2012; 45(2):129-35.
16. Sirtes G, Waltimo T, Schaetzle M, Zehnder M. The Effects of Temperature on Sodium Hypochlorite Short-Term Stability, Pulp Dissolution Capacity, and Antimicrobial Efficacy. *J Endod.* 2005; 31(9):669-71.
  17. Rossi-Fedele G, De Figueiredo JAP. Use of a bottle warmer to increase 4% sodium hypochlorite tissue dissolution ability on bovine pulp. *Aust Endod J.* 2008; 34(1):39-42.
  18. Clarkson RM, Kidd B, Evans GE, Moule AJ. The Effect of Surfactant on the Dissolution of Porcine Pulpal Tissue by Sodium Hypochlorite Solutions. *J Endod.* 2012; 38(9):1257-60.
  19. Almeida LHS de, Leonardo NG e S, Gomes APN, Giardino L, Souza EM, Pappen FG. Pulp Tissue Dissolution Capacity of Sodium Hypochlorite Combined with Cetrimide and Polypropylene Glycol. *Braz Dent J.* 2013; 24(5):477-81.
  20. Slutzky-Goldberg I, Hanut A, Matalon S, Baev V, Slutzky H. The Effect of Dentin on the Pulp Tissue Dissolution Capacity of Sodium Hypochlorite and Calcium Hydroxide. *J Endod.* 2013; 39(8):980-3.
  21. De Almeida LHS, Leonardo NG e S, Gomes APN, Souza EM, Pappen FG. Influence of EDTA and dentine in tissue dissolution ability of sodium hypochlorite: Pulp Dissolution of Sodium Hypochlorite. *Aust Endod J.* 2013.
  22. Echeverri D, Acuña C. Dissolution of Connective Tissue in Sodium Hypochlorite Alone and in Combination with 3% Hydrogen Peroxide. *Int J Odontostomatol.* 2012; 6(3):263-6.
  23. Al-Jadaa A, Paqué F, Attin T, Zehnder M. Necrotic pulp tissue dissolution by passive ultrasonic irrigation in simulated accessory canals: impact of canal location and angulation. *Int Endod J.* 2009; 42(1):59-65.
  24. Kuhn K, Rudolph H, Luthardt RG, Stock K, Diebolder R, Hibst R. Er:YAG Laser Activation of Sodium Hypochlorite for Root Canal Soft Tissue Dissolution: Er:YAG/ NaOCl Root Canal Tissue Dissolution. *Lasers Surg Med.* 2013; 45(5):339-44.
  25. Guneser MB, Arslan D, Usumez A. Tissue dissolution ability of sodium hypochlorite activated by photon-initiated photoacoustic streaming technique. *J Endod.* 2015; 41(5):729-32.
  26. Rossi-Fedele G, Steier L, Dogramaci EJ, Canullo L, Steier G, de Figueiredo JAP. Bovine pulp tissue dissolution ability of HealOzone®, Aquatone Alpha Electrolyte® and sodium hypochlorite: Ozone, EAS and NaOCl Pulp Dissolution Ability. *Aust Endod J.* 2013; 39(2):57-61.
  27. Yang S, Rivera E, Baumgardner K, Walton R, Stanford C. Anaerobic tissue-dissolving abilities of calcium hydroxide and sodium hypochlorite. *J Endod.* 1995; 21(12):613-6.
  28. Okino LA, Siqueira EL, Santos M, Bombana AC, Figueiredo JAP. Dissolution of pulp tissue by aqueous solution of chlorhexidine digluconate and chlorhexidine digluconate gel. *Int Endod J.* 2004; 37(1):38-41.
  29. Khademi A, Usefian E, Feizianfard M. Tissue dissolving ability of several endodontic irrigants on bovine pulp tissue. *Iran Endod J.* 2007; 2(2):65-8.
  30. Couto De Oliveira G, Ferraz CS, Andrade Júnior CV, Pithon MM. Chlorhexidine gel associated with papain in pulp tissue dissolution. *Restor Dent Endod.* 2013; 38(4):210
  31. Beltz RE, Torabinejad M, Pouresmail M. Quantitative analysis of the solubilizing action of MTAD, sodium hypochlorite, and EDTA on bovine pulp and dentin. *J Endod.* 2003; 29(5):334-7.
  32. Ballal NV, Mala K, Bhat KS. Effect of maleic acid and ethylenediaminetetraacetic acid on the dissolution of human pulp tissue - an *in vitro* study: Pulp-dissolution effect of Maleic acid and EDTA. *Int Endod J.* 2011; 44(4):353-6.
  33. Irala LED, Grazziotin-Soares R, Salles AA, Munari AZ, Pereira JS. Dissolution of bovine pulp tissue in solutions consisting of varying NaOCl concentrations and combined with EDTA. *Braz Oral Res.* 2010; 24(3):271-6.
  34. Cobankara FK, Ozkan HB, Terlemez A. Comparison of Organic Tissue Dissolution Capacities of Sodium Hypochlorite and Chlorine Dioxide. *J Endod.* 2010; 36(2):272-4.
  35. Singh S, Sinha R, Kar SK, Ather A, Limaye SN. Effect of chlorine dioxide and sodium hypochlorite on the dissolution of human pulp tissue – An *in vitro* study. *Med J Armed Forces India.* 2012; 68(4):356-9.
  36. Taneja S, Mishra N, Malik S. Comparative evaluation of human pulp tissue dissolution by different concentrations of chlorine dioxide, calcium hypochlorite and sodium hypochlorite: An *in vitro* study. *J Conserv Dent.* 2014; 17:541-5
  37. Jain A, Shrivastava TV, Tabassum S, Bahuguna R. Comparison of human pulp tissue dissolution capacities of different irrigating solutions: An *in vitro* study. *Eur J Gen Dent.* 2015; 4:64-7.
  38. Dutta A, Saunders WP. Comparative evaluation of calcium hypochlorite and sodium hypochlorite on soft-tissue dissolution. *J Endod.* 2012; 38(10):1395-8.
  39. Arslan D, Guneser MB, Kustarci A, Er K, Siso SH. Pulp tissue dissolution capacity of QMix 2in1 irrigation solution. *Eur J Dent.* 2015; 9(3):423-7.
  40. Rao SA, Sunitha L, Rao BN, Naik JP, Shekar VC. Efficacy of garlic extract and sodium hypochlorite on dental pulp dissolution: An *in vitro* study. *Saudi Endod J.* 2017; 7:36-9.